

1.13 Function Model Selection

AP Precalculus

1.13 Practice

Select the appropriate model for the data (linear/quadratic/cubic). Explain why it models the data.

1.

x	1	2	3	4	5	6
y	4	15	50	121	240	419

$11, 35, 71, 119, 179$
 $24, 36, 48, 60$
 $12, 12, 12$

Cubic, third difference constant of 12

2.

x	1	2	3	4	5	6
y	49	46	43	40	37	34

$-3, -3, -3, -3, -3$

Linear, first difference constant of -3.

3.

x	1	2	3	4	5	6
y	119	110	95	74	47	14

$9, 15, 21, 27, 33$
 $6, 6, 6, 6$

Quadratic, second difference constant of 6

4.

x	2	4	5	6	7	8
y	23	81	122	171	228	293

$58, 90, 122$
 $32, 32$

Quadratic, second difference constant of 32

CALCULATOR ACTIVE. Use the model to answer the questions in context.

5. A patient receives a dose of painkiller. The function $p(t) = \frac{2t^2 + 10t}{t^3 + 1}$ models the amount of painkiller in the blood stream over time, where t is time in hours and p is painkiller in milligrams.

- a. Find $p(2)$. Explain your solution in context.

$$f(2) = 3.111$$

Two hours after taking the painkiller, there was 3.111 mg of painkiller in the blood stream.

- b. What is the average rate of change from $t = 1$ to $t = 2$? Explain your solution in context.

$$\frac{6 - 3.111}{1 - 2} = -2.889 \quad \text{The painkiller is leaving the blood stream at -2.889 mg per hour.}$$

- c. What is the maximum amount of painkiller in the patient's bloodstream?

$$6.161 \text{ mg}$$

6. A rectangle is inscribed in a circle with diameter of 12 cm. The width of the rectangle is x cm. The function $A(x) = x\sqrt{144 - x^2}$ models the area of the rectangle.

- a. What is the restricted domain of the function?


$$0 \leq x \leq 12$$

- b. What is the restricted range of the function?

$$0 \leq y \leq 72$$

- c. What is the maximum area of the rectangle?

$$72 \text{ cm}^2$$



Use the graph of the piecewise function to answer the questions in context.

7. Generic Strawberry Fields allows customers to pay \$5 to pick strawberries plus 50 cents for every pint or partial pint of strawberries that they pick. There is a limit of 8 pints per customer. The piecewise function f shown below models the price of strawberries picked.

- a. What is the domain in this context?


$$0 \leq x \leq 8 \text{ pints of strawberries}$$

- b. The range can be represented by $y = 5 + 0.5x$ where x is integers in the domain. Explain why.

The range starts at 5 and increases in increments of 0.50. Using only integers for the domain generates the range of 5, 5.5, 6, 6.5, 7, 7.5, 8, 8.5, 9

- c. Find $f(3.5)$. What does it mean in this context?

$$f(3.5) = 7 \text{ 3.5 pints of strawberries cost 7 dollars.}$$



8. The piecewise function f shown below models the insulin levels of a patient over time where $x = 0$ represents 8:00.

- a. What is the domain in this context?

$$0 \leq x \leq 10 \text{ hours after 8:00}$$

- b. What is the range in this context?

$$\text{This patient's insulin levels are } 8 \leq y \leq 24$$

Possible insulin levels are $0 \leq y \leq \text{around } 35$ for most patients.


- c. Find $f(3)$. What does it mean in this context?

$$f(3) = 24 \text{ 3 hours after 8:00 (or 11:00), the patient's insulin level is 24 mIU/ml.}$$

- d. Find average rate of change from $[0, 3]$. Explain the meaning in this context.

$$\frac{24-8}{3-0} = \frac{16}{3} = 5.333$$

The patient's insulin level increased 5.333 levels every hour for the first 3 hours.



1.13 Function Model Selection

1.13 Test Prep

Multiple Choice - CALCULATOR ACTIVE

For questions 9 and 10, use the table of values for polynomial f at selected values of x .

x	2	4	6	8	10	12
$f(x)$	97	405	977	1861	3105	4757


9. Which of the following statements are true?

- (A) The function is quadratic because second differences are a nonzero constant
- (B) The function is quadratic because third differences are a nonzero constant
- (C) The function is cubic because second differences are a nonzero constant
- (D) The function is cubic because third differences are a nonzero constant

10. The average rate of change from $x = -22$ to $x = 4$ is 34. Which statement best describes the $f(x)$?

- (A) There must be at least one zero on the interval $[-22, 4]$ because $f(-22)$ is negative.
- (B) The number of zeros on the interval $[-22, 4]$ cannot be determined because $f(-22)$ is positive.
- (C) There must be at least one zero on the interval $[-22, 4]$ because the average rate of change is negative.
- (D) There is at least one zero in that interval $[-22, 4]$ because the average rate of change is positive.

11. A track athlete is running laps. The graph shows the average rate of change from the previous lap.



A function model T is constructed for the time of each lap. Which of the following statements best supports the selection of the model of a model for T ?

- (A) Since the rate of change is roughly linear, a linear model is best for T .
- (B) Since the rate of change is roughly linear, a quadratic model is best for T .
- (C) Since the rate of change is roughly linear, a cubic model is best for T .
- (D) Since the rate of change is negative and positive, a quadratic model is best for T .