

### 3.15 Rates of Change in Polar Functions

AP Precalculus

### 3.15 Practice

Use the table of selected values for the polar function  $r = f(\theta)$  to answer the following.

1.

$\theta$	0	$\frac{\pi}{4}$	$\frac{\pi}{2}$	$\frac{3\pi}{4}$	$\pi$	$\frac{5\pi}{4}$	$\frac{3\pi}{2}$	$\frac{7\pi}{4}$	$2\pi$
$r$	11	9.53	6	2.46	1	2.46	6	9.53	11

a. Determine the interval(s) where  $f$  is increasing. Determine the interval(s) where  $f$  is decreasing.

increasing  $(\pi, 2\pi)$

decreasing  $(0, \pi)$

b. The distance between  $f(\theta)$  and the pole is increasing or decreasing on the interval  $\pi \leq \theta \leq 2\pi$ . Justify your answer.

$r$  is positive and increasing so the distance is increasing

c. Find the average rate of change of  $f$  between  $\theta = \frac{5\pi}{4}$  and  $\theta = \frac{7\pi}{4}$ .

$$\frac{9.53 - 2.46}{\frac{7\pi}{4} - \frac{5\pi}{4}} = \frac{7.07}{\frac{2\pi}{4}} = \frac{7.07}{\frac{\pi}{2}} = \frac{14.14}{\pi} \approx 4.5 \text{ units per radian}$$

d. Estimate the value of  $f\left(\frac{\pi}{3}\right)$  using an average rate of change.

Use the interval  $\left[\frac{\pi}{4}, \frac{\pi}{2}\right]$

$$\frac{6 - 9.53}{\frac{\pi}{2} - \frac{\pi}{4}} = \frac{-3.53}{\frac{\pi}{4}} = \frac{-14.16}{\pi} \approx -4.495$$

$$y - y_1 = m(x - x_1)$$

$$y - 6 = -4.495(x - \frac{\pi}{2})$$

$$y - 6 = -4.495x + 7.06$$

$$+6 \quad +6$$

$$y = -4.495x + 13.06$$

Are there any extrema on the interval  $\left[\frac{\pi}{4}, \frac{5\pi}{4}\right]$ ? Explain how you know.

yes, there is at least one because the function goes from decreasing to increasing

$$\boxed{\frac{\pi}{3} \approx 1.05}$$

2.

a. Is  $f$  increasing or decreasing on the interval  $\frac{\pi}{4} \leq \theta \leq \frac{\pi}{2}$ ?

increasing

b. Is the distance between  $f(\theta)$  and the pole is increasing or decreasing on the interval  $0 \leq \theta \leq \frac{\pi}{4}$ ?

$r$  is negative and decreasing so the distance is increasing

c. Is the rate of change of  $f$  faster on the interval  $\left[0, \frac{\pi}{8}\right]$  or the interval  $\left[\frac{\pi}{8}, \frac{\pi}{4}\right]$ ? Justify.

$$\frac{-1.41 - 0}{\frac{\pi}{8} - 0} = \frac{-1.41}{\frac{\pi}{8}} = \frac{-11.28}{\pi} \approx -3.59$$

$$\frac{-2 - (-1.41)}{\frac{\pi}{4} - \frac{\pi}{8}} = \frac{-0.59}{\frac{\pi}{8}} = \frac{-4.72}{\pi} \approx -1.502$$

Faster on  $\left[0, \frac{\pi}{8}\right]$  because  $-3.59$  is more negative (steeper) than  $-1.502$

$\theta$	$r$
0	0
$\frac{\pi}{8}$	-1.41
$\frac{\pi}{4}$	-2
$\frac{3\pi}{8}$	-1.41
$\frac{\pi}{2}$	0

Use the polar function  $r = f(\theta)$  to fill in the table and answer the questions. Calculator Active.

3.  $r = f(\theta) = 8 \cos(\theta)$

$\frac{2\pi}{3}$

$\theta$	0	$\frac{\pi}{6}$	$\frac{\pi}{2}$	$\frac{5\pi}{6}$	$\pi$	$\frac{7\pi}{6}$	$\frac{3\pi}{2}$	$\frac{11\pi}{6}$	$2\pi$
$r$	8	6.928	0	-6.928	-8	-6.928	0	6.928	8

a. Determine the interval(s) where  $f$  is increasing. Determine the interval(s) where  $f$  is decreasing.

increasing on  $(\pi, 2\pi)$

decreasing on  $(0, \pi)$

b. How many extrema on the interval  $\frac{5\pi}{6} \leq \theta \leq \frac{11\pi}{6}$ ? Justify.

there is at least one because the function goes from decreasing to increasing

c. Determine the intervals where the distance between  $f(\theta)$  and the pole is increasing on the interval  $0 \leq \theta \leq 2\pi$ . Justify your answer.

increasing on  $(\frac{\pi}{2}, \pi)$  because  $r$  is negative and decreasing

increasing on  $(\frac{3\pi}{2}, 2\pi)$  because  $r$  is positive and increasing

d. Determine the intervals where the distance between  $f(\theta)$  and the pole is decreasing on the interval  $0 \leq \theta \leq 2\pi$ . Justify your answer.

decreasing on  $(\pi, \frac{3\pi}{2})$  because  $r$  is negative and increasing

decreasing on  $(0, \frac{\pi}{2})$  because  $r$  is positive and decreasing

e. Find the average rate of change of  $f$  between  $\theta = \frac{\pi}{2}$  and  $\theta = \frac{5\pi}{6}$ . Use to estimate  $f(\frac{2\pi}{3})$ .

$$\frac{0 - (-6.928)}{\frac{\pi}{2} - \frac{5\pi}{6}} = \frac{6.928}{-\frac{2\pi}{6}} = \frac{6.928}{-\frac{\pi}{3}} = \frac{6.928}{-\pi} \approx -6.615$$

$$y - y_1 = m(x - x_1)$$

$$y - 0 = -6.615(x - \frac{\pi}{2})$$

$$y = -6.615x + 10.39$$

$$y = -6.615(\frac{2\pi}{3}) + \square$$

$$y = -3.464$$

$$f(\frac{2\pi}{3}) \approx -3.464$$

4.  $r = f(\theta) = -3 + 5 \sin(\theta)$

- a. Is the distance between  $f(\theta)$  and the pole is increasing or decreasing on the interval  $\frac{\pi}{2} \leq \theta \leq \frac{3\pi}{4}$ ?

decreasing on  $(\frac{\pi}{2}, \frac{3\pi}{4})$  because  $r$  is positive and decreasing

- b. Find the average rate of change of  $f$  between  $\theta = \frac{\pi}{4}$  and  $\theta = \frac{\pi}{2}$ .

$$\frac{2 - 0.535}{\frac{\pi}{2} - \frac{\pi}{4}} = \frac{1.465}{\frac{\pi}{4}} = \frac{5.86}{\pi} \approx 1.865 \text{ units per radian}$$

- c. Estimate the value of  $f(\frac{5\pi}{6})$  using an average rate of change

Use the interval  $[\frac{3\pi}{4}, \pi]$   $\frac{0.535 - (-3)}{\frac{3\pi}{4} - \pi} = \frac{3.535}{-\frac{\pi}{4}} = \frac{14.14}{-\pi} \approx -4.5$

$$y - (-3) = -4.5(x - \pi)$$

$$\boxed{\phantom{000}} = -4.5x + 4.5\pi$$

$$\phantom{\boxed{\phantom{000}}} - 3$$

$$y = -4.5x + 11.137$$

$$\boxed{\phantom{000}} \quad 4.5\left(\frac{5\pi}{6}\right) + 11.137 = -0.643$$

$$f\left(\frac{5\pi}{6}\right) \approx -0.643$$

$\theta$	$r$
0	-3
$\frac{\pi}{4}$	0.535
$\frac{\pi}{2}$	2
$\frac{3\pi}{4}$	0.535
$\pi$	-3

### 3.15 Rates of Change in Polar Functions

### 3.15 Test Prep

5. Consider the graph of the polar function  $r = f(\theta)$ , where  $\theta$  is increasing in the polar coordinate system on the interval  $0 \leq \theta \leq 2\pi$ . Given  $f(\theta) < 0$  and decreasing on the interval  $\pi \leq \theta \leq \frac{3\pi}{2}$  which of the following statements is true about the distance between the point with polar coordinates  $(f(\theta), \theta)$  and the origin.

- (A) The distance is increasing for  $0 \leq \theta \leq 2\pi$ .
- (B) The distance is decreasing for  $0 \leq \theta \leq 2\pi$ .
- (C)** The distance is increasing for  $\pi \leq \theta \leq \frac{3\pi}{2}$ .
- (D) The distance is decreasing for  $\pi \leq \theta \leq \frac{3\pi}{2}$ .

because  $r$  is negative and decreasing on  $(\pi, \frac{3\pi}{2})$  so the distance is increasing

Use the table of selected values for the polar equation  $r = f(\theta)$  below to answer questions 6 and 7.

CALCULATOR ACTIVE

$\theta$	0	$\frac{\pi}{8}$	$\frac{\pi}{4}$
$r$	-1	$-\frac{\sqrt{3}}{2}$	$-\frac{1}{2}$

$-1 \quad -0.866 \quad -0.5$

$r$  is negative and increasing so the distance is decreasing

6. The graph of the polar function  $r = f(\theta)$ , is given the polar coordinate system. Which of the following descriptions is true?

- (A) As  $\theta$  increasing from 0 to  $\frac{\pi}{4}$ , the polar function  $r = f(\theta)$  is increasing, and the distance between the point  $(f(\theta), \theta)$  on the curve and the origin is increasing.
- (B) As  $\theta$  increasing from 0 to  $\frac{\pi}{4}$ , the polar function  $r = f(\theta)$  is increasing, and the distance between the point  $(f(\theta), \theta)$  on the curve and the origin is decreasing.
- (C) As  $\theta$  increasing from 0 to  $\frac{\pi}{4}$ , the polar function  $r = f(\theta)$  is decreasing, and the distance between the point  $(f(\theta), \theta)$  on the curve and the origin is increasing.
- (D) As  $\theta$  increasing from 0 to  $\frac{\pi}{4}$ , the polar function  $r = f(\theta)$  is decreasing, and the distance between the point  $(f(\theta), \theta)$  on the curve and the origin is decreasing.

=

7. If the value of  $r = f\left(\frac{\pi}{8}\right)$  is estimated using the average rate of change of the function over the interval  $0 \leq \theta \leq \frac{\pi}{4}$ , which of the following is true?

- (A) The estimated value would be an overestimate of the actual value by approximately 0.116.
- (B) The estimated value would be an underestimate of the actual value by approximately  $-0.884$ .
- (C) The estimated value would be an overestimate of the actual value by approximately 1.616.
- (D) The estimated value would be an underestimate of the actual value by approximately  $-1.043$ .

Use the interval  $\left[0, \frac{\pi}{4}\right]$  to estimate

$$\frac{-0.5 - (-1)}{\frac{\pi}{4} - 0} = \frac{0.5}{\frac{\pi}{4}} = \frac{2}{\pi} \approx 0.636$$

$$y - (-1) = \frac{2}{\pi}(x - 0)$$

$$y + 1 = \frac{2}{\pi}x$$

$$y = \frac{2}{\pi}x - 1$$

$$y = \frac{2}{\pi}\left(\frac{\pi}{8}\right) - 1$$

$$y = \frac{2}{8} - 1 = -0.75$$

actual  $\rightarrow f\left(\frac{\pi}{8}\right) = -\frac{\sqrt{3}}{2} \approx -0.866$   
 estimate  $\rightarrow f\left(\frac{\pi}{8}\right) \approx -0.75$

The estimate of  $-0.75$  is above the actual value  $-0.866$  by 0.116.